

Luminous hot accretion flows: thermal equilibrium curve and thermal stability

Feng Yuan

*Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138;
fyuan@cfa.harvard.edu*

ABSTRACT

In a previous paper, we presented the global solution of a new accretion flow model, namely luminous hot accretion flows (LHAFs). In this *Letter*, we first show the corresponding thermal equilibrium curve of LHAFs in the mass accretion rate vs. surface density diagram. Then we examine its thermal stability against local perturbations. We find that LHAFs are thermally unstable when thermal conduction is neglected. However, when the accretion rate is not very large, the timescale of the growth of perturbations is longer than the accretion timescale, therefore the instability has no dynamical effect on the accretion flow. When the accretion rate is large, the perturbations can grow very fast at a certain radius. As a result, some cold clumps may form and the accretion flow will become multi-phase.

Subject headings: accretion, accretion disks — black hole physics — galaxies: active — galaxies: nuclei

1. Introduction: the physics of the new hot accretion disk solution

There has been a great interest to the accretion process around black holes. The most famous accretion solution is the geometrically thin and optically thick cold disk model developed by Shakura & Sunyaev (1973; hereafter SSD) and others. The second solution was discovered by Shapiro, Lightman & Eardley (1976; hereafter SLE). This solution is optically thin and hot, with $T_e \sim 10^9 \text{K}$. Pringle (1976) found that SLE is thermally unstable although it is not clear what is the consequence of such instability.

In both SSD and SLE, the energy advection is neglected. It was found that when the mass accretion rate is higher than the Eddington rate $\dot{M}_{\text{Edd}} (\equiv 10L_{\text{Edd}}/c^2)$, the large optical depth traps most of the photons therefore most of the viscously dissipated energy is stored in

the gas and advected into the black hole rather than radiated away (“slim disk”; Abramowicz et al. 1988).

The fourth accretion solution is the advection-dominated accretion flow (ADAF; Narayan & Yi 1995; Abramowicz et al. 1995, hereafter A95; see reviews by Narayan, Mahadevan & Quataert 1998; Kato, Fukue & Mineshige 1998). Different from SLE, the energy advection is included in the ions energy equation of an ADAF, $Q_{\text{adv}} = Q_{\text{vis}} - Q_{\text{ie}}$. Here Q_{adv} , Q_{vis} and Q_{ie} are the rates of energy advection, viscous heating and Coulomb cooling per unit area of the accretion disk, respectively. In a typical ADAF, the density of gas is very low, so $Q_{\text{ie}} \ll Q_{\text{vis}} \approx Q_{\text{adv}}$, i.e., *the viscous heating is balanced by advective cooling*. Since $Q_{\text{ie}} \propto \dot{M}^2$ while $Q_{\text{vis}} \propto \dot{M}$, i.e., Q_{ie} increases faster than Q_{vis} with increasing $\dot{M}$, there exists a critical rate $\dot{M}_1$, determined by $Q_{\text{vis}} \approx Q_{\text{ie}}$. At this rate, a large fraction of the viscously dissipated energy is transferred to the electrons and radiated away, so the accretion flow ceases to be advection-dominated. Above $\dot{M}_1$, it was thought previously that no *hot* accretion solution exists and the only viable solution is SSD.

However, our recent work (Yuan 2001, hereafter Y01) indicated that this is not true: above $\dot{M}_1$, a new hot accretion solution exists. To illustrate, let’s first write out the formula of the energy advection of ions,

$$Q_{\text{adv}} = \rho H v_r \left[\frac{k}{\mu m_\mu} \frac{1}{\gamma - 1} \frac{dT}{dr} - \frac{kT}{\mu m_\mu} \frac{1}{\rho} \frac{d\rho}{dr} \right] \equiv Q_{\text{int}} - Q_{\text{com}}, \quad (1)$$

i.e., the energy advection consists of two terms, namely the internal energy gradient term Q_{int} and the compression work Q_{com} . *The full condition for the existence of a hot accretion solution, when the flow starts out hot, is $dT/dr < 0$, i.e., $Q_{\text{int}} > 0$ (note $v_r < 0$).* For an ADAF, $\dot{M} < \dot{M}_1$, $Q_{\text{adv}} = Q_{\text{vis}} - Q_{\text{ie}} > 0$, so $Q_{\text{int}} = Q_{\text{adv}} + Q_{\text{com}} > 0$. When $\dot{M} > \dot{M}_1$, $Q_{\text{adv}} = Q_{\text{vis}} - Q_{\text{ie}} < 0$, but there obviously exists another critical accretion rate, $\dot{M}_2$, determined by $Q_{\text{com}} + Q_{\text{vis}} \approx Q_{\text{ie}}$. Below $\dot{M}_2$, we still have $Q_{\text{int}} = Q_{\text{vis}} + Q_{\text{com}} - Q_{\text{ie}} > 0$. Therefore, the accretion flow can remain hot if it starts out hot. We denote this new solution as luminous hot accretion flow (LHAF hereafter).

From the above analysis, we see that LHAFs are along the line of ADAFs—the equations describing both are completely the same and we just extend ADAFs to higher accretion rates. However, the dynamics of an LHAF is quite different from an ADAF. In an ADAF, $Q_{\text{adv}} > 0$, i.e., the advection plays a “cooling” role from the Lagrangian point of view. But in an LHAF, $Q_{\text{adv}} < 0$, so it plays a “heating” role. In the language of entropy, it is the conversion of entropy together with viscous dissipation that supplies the radiation of LHAFs. In this sense, an LHAF is dynamically similar with Bondi accretion and cooling flow in galactic clusters.

In this context we can understand why previous authors didn’t find this solution. In Narayan & Yi (1995) and Esin et al. (1997), they *a priori* set the advection factor “ f ” ($\equiv$

$Q_{\text{adv}}/Q_{\text{vis}}$) as positive, so they only obtained the ADAF solution. Similarly, A95 and Chen et al. (1995, hereafter C95) didn't find this solution because they set the parameter ξ (see eq. (6) below) as positive.

In this *Letter*, we will first show the thermal equilibria of LHAFs in the $\dot{M}$ vs. Σ (surface density) diagram (§2). Such a diagram is widely used in the study of accretion disks (e.g., A95; C95; Kusunose & Mineshige 1996; Björnsson et al. 1996). Then in §3 we investigate the thermal stability of LHAFs.

2. Thermal equilibrium curve of LHAFs

We take a one-temperature accretion flow as an example for simplicity. For the purpose of comparison, we adopt almost exactly the same equations as C95 (see also A95).

$$\dot{M} = -2\pi R \Sigma v_r, \quad (2)$$

$$\nu \Sigma = \frac{\dot{M}}{3\pi} f_* g^{-1}, \quad (3)$$

$$Q_{\text{adv}} = Q_{\text{vis}} - Q_{\text{rad}}. \quad (4)$$

They represent conservations of mass, angular momentum, and energy, respectively. Here $\nu = \frac{2}{3}\alpha c_s H$ is the kinetic viscosity coefficient, $\Sigma = 2\rho H$ is the surface density, $f_* = 1 - 9\Omega(3R_s)/[\Omega(R)(R/R_s)^2]$ with $R_s = 2GM/c^2$, $g = -\frac{2}{3}(d\ln\Omega/d\ln R)$. The forms of Q_{vis} , Q_{adv} and Q_{rad} are,

$$Q_{\text{vis}} = \frac{3\dot{M}}{4\pi} \Omega^2 f_* g, \quad (5)$$

$$Q_{\text{adv}} = -\Sigma v_r \frac{p}{\rho} \xi = \frac{\dot{M}}{2\pi R^2} \frac{p}{\rho} \xi, \quad (6)$$

$$Q_{\text{rad}} = 8\sigma T^4 \left(\frac{3\tau}{2} + \sqrt{3} + \frac{8\sigma T^4}{Q_{\text{brem}}} \right)^{-1}. \quad (7)$$

The optical depth $\tau = \tau_{\text{es}} + \tau_{\text{abs}}$, with $\tau_{\text{es}} = (1/2)k_{\text{es}}\Sigma$ and $\tau_{\text{abs}} = Q_{\text{brem}}/(8\sigma T^4)$. The equation of state is $p = p_{\text{gas}} + p_{\text{rad}}$, with $p_{\text{gas}} = \frac{\mathfrak{R}}{\mu}\rho T$, and $p_{\text{rad}} = \frac{Q_{\text{rad}}}{4c} \left(\tau + \frac{2}{\sqrt{3}} \right)$ (Abramowicz et al. 1996). The bremsstrahlung radiation $Q_{\text{brem}} = 2H(q_{ei} + q_{ee})$. The formula for q_{ei} and q_{ee} are from Narayan & Yi (1995). Another useful equation is the hydrostatic balance equation, $H = c_s/\Omega_K$, with $c_s = (p/\rho)^{1/2}$ is the local sound speed. The Paczyński & Wiita (1980) potential is used.

We solve eqs. (2)-(7) to obtain the thermal equilibrium curve of LHAFs. One noticeable parameter is ξ in eq. (6). From §1, we know that for an LHAF, $Q_{\text{adv}} < 0$, therefore we should set $\xi < 0$ to recover this solution. We can recover all other accretion solutions by simply setting $\xi = 1$, as in A95. The crudeness of the value of ξ doesn't affect our qualitative results although it does prevent us from obtaining the exact quantitative results such as the ranges of accretion rates to which various accretion disk models correspond.

Figure 1 shows the thermal equilibrium curves of different solutions. Comparing with the figures in A95 or C95, we see that the three “U” shaped thick lines are new and they denote LHAFs with different values of ξ . These lines extend from optically thin regime to optically thick regime, bridging the SLE and the SSD solutions. It is not clear what physical implications the “bridging” has, or it might just be a mathematical trick. But we note in this context that in the slim disk solutions, they did find solutions with the advection term being negative in some cases (Abramowicz et al. 1988; Szuszkiewicz, private communication). Some segment of the “U” curves are superimposed on the SLE and SSD lines. We find that advection in these “superimposed” segment is equal to zero, same with SLE and SSD. Only the “separate” segment denotes the genuine LHAFs, since for this segment the energy advection is negative and plays a significant role in the energy balance of ions. We see that above the critical accretion rate of ADAF, $\dot{M}_1 (\sim 0.1 \dot{M}_{\text{Edd}})$, both ADAF and SLE disappear and LHAF is the only available *hot* solution. This is consistent with our analysis in §1.

For comparison, we also solve the *global* solutions of the standard accretion equations describing a one-temperature accretion flow with the same parameters as in Fig. 1 (see Yuan 1999 for the equations except that the radiative term is now replaced by eq. (7) of the present paper). The results are denoted by the filled circles (for ADAFs) and triangles (for LHAFs). We can see that our local algebraic analysis is qualitatively consistent with the global results, i.e., when the accretion rate is lower than a certain value, the solutions are ADAFs; while above this value, the solutions are LHAFs. We also calculated the value of ξ for a specific example of $\dot{M} = 10^{-1.6} \dot{M}_{\text{Edd}}$ and the outer boundary conditions of $R_{\text{out}} = 10^3 R_s$, $T_{\text{out}} = 2 \times 10^9 \text{K}$, and $v/c_s = 0.4$. We obtained $\xi \approx -0.3$ at $R = 5R_s$.

We want to emphasize that our “U” shaped line is completely different from the line with the similar shape in C95, which was obtained by setting a very large viscous parameter $\alpha > \alpha_{cr}$ with $\alpha_{cr} \sim 0.2$ or larger. In fact, a later more accurate treatment of microphysics and the inner boundary condition by Björnsson et al. (1996) indicated that $\alpha_{cr} > 1$, therefore, the “U” shaped branch in C95 is unphysical.

3. Thermal stability

From the density profile of the global solution in Y01, we know that LHAFs are viscously stable. An important problem then is to analyze its thermal stability. Before doing that, we note that an LHAF is dynamically very similar to a cooling flow in galactic clusters. Many authors have studied the thermal stability of cooling flow and concluded that it is unstable against local perturbations if thermal conduction is neglected (e.g., Fabian & Nulsen 1977; Mathews & Bregman 1978; Nulsen 1986). Thermal conduction can strongly stabilize the cooling flow (Zakamska & Narayan 2003; Kim & Narayan 2003).

Now let's investigate the thermal stability of LHAFs against local perturbations. Such perturbations can be, e.g., that the local gas density increases wherever the initial ambient magnetic field is less than average. We follow the standard analysis approach presented by Kato, Abramowicz & Chen (1996). Taking the surface density as an example, we denote perturbations Σ_1 superimposed over the unperturbed quantities Σ as $\sigma \equiv \Sigma_1/\Sigma \propto \exp(n\Omega t - ikr)$ where $n\Omega$ is the growth rate of perturbation and k is the perturbation wave number. Substituting the perturbed quantities to the time-dependent accretion equations, we can obtain the perturbed equations and the dispersion relation. The arguments for ADAFs presented in Kato et al. (1996) hold for LHAFs as well although the advection is negative and radiative cooling is important here. We assume that thermal conductivity is suppressed by the presence of the tangled magnetic field in the accretion flow (but see Narayan & Medvedev 2001). We expect the flow will be thermally stable if thermal conduction is strong. The dispersion relation is,

$$-3n\Omega\sigma = G\sigma, \quad (8)$$

and the condition for instability is

$$G \equiv -\frac{Q_{\text{vis}}}{W} \left(\frac{\partial \ln \eta}{\partial \ln \Sigma} \right)_W - \frac{Q_{\text{rad}}}{W} \left(\frac{\partial \ln Q_{\text{rad}}}{\partial \ln \Sigma} \right)_W < 0. \quad (9)$$

Here $W \equiv 2Hp$ and $\eta \equiv \nu\Sigma$. Since $\eta \propto \Sigma c_s^2/\Omega \propto W/\Omega$, so $(\partial \ln \eta / \partial \ln \Sigma)_W = 0$. For a one-temperature LHAF, $Q_{\text{rad}} \propto H\rho^2 T^{1/2} \propto \Sigma^2 T^{1/2}/H \propto \Sigma^2$, so

$$(\partial \ln Q_{\text{rad}} / \partial \ln \Sigma)_W > 0. \quad (10)$$

Therefore one-temperature LHAFs are thermally unstable against local perturbations.

The case of two-temperature LHAF is a little bit more complicated. Treating the energy equation in Kato et al. (1996) as the energy equation of ions, the instability condition is exactly the same as eq. (9) except that we should replace Q_{rad} with Q_{ie} , the Coulomb collision cooling rate. Since $Q_{ie} \propto \rho^2 H T_e^{-3/2} T_i$, to determine the sign of G , we must turn to the electron energy equation to obtain the relation between T_e and T_i . Writing the energy

equation of electrons as $Q_{\text{adv,e}} = Q_{ie} - Q_{\text{rad,e}}$, with $Q_{\text{adv,e}}$ and $Q_{\text{rad,e}}$ are the energy advection and radiative cooling of electrons respectively, the global solution of Y01 indicates that for LHAF a good approximation to the electrons energy equation is $Q_{ie} \approx Q_{\text{adv,e}}$ in the region outside of $\sim 100R_s$ and $Q_{ie} \approx Q_{\text{rad,e}}$ inside $100R_s$ (see Figure 5 in Y01). Taking $Q_{ie} \approx Q_{\text{adv,e}}$ we can get $T_e \propto W^{1/5}\Sigma^{1/5}$. Hence $Q_{ie} \propto \Sigma^{6/5}W^{1/5}$ and

$$G = -\frac{Q_{ie}}{W} \left(\frac{\partial \ln Q_{ie}}{\partial \ln \Sigma} \right)_W = \frac{6}{5} > 0. \quad (11)$$

So a two-temperature LHAF is thermally unstable at least at the region outside of $\sim 100R_s$. Inside $\sim 100R_s$, the radiative cooling is in general dominated by the thermal Comptonization. The seed photons can be the synchrotron photons, or other soft photons from cold matter such as a SSD underlying or outside of the LHAF, or cold clumps suspending in the hot gas. The origin of the cold clumps in LHAFs can be due to the thermal instability of LHAFs (see below), or originally existed in the accretion material. For luminous sources the soft photons from the cold matter is likely to dominate over the synchrotron photons as the seed photons of Comptonization (e.g., Zdziarski et al. 1998). Since $Q_{\text{rad,e}} = (4kT_e/m_e c^2)\rho H\kappa_{es}U_r c$, with U_r denotes the flux of seed photon, using $Q_{ie} \approx Q_{\text{rad,e}}$, we find that if $U_r \propto T_i^2$ or steeper, the two-temperature LHAFs will be thermally stable, otherwise it is unstable. Unfortunately, it is not easy to determine whether this condition can be satisfied or not, because of the complicated energy interaction between the cold and hot phases in accretion flow (e.g., Ferland & Rees 1988).

Now let's discuss the consequence of the thermal instability against local perturbations. An important factor is how fast the perturbations grow compared to other related timescales. It is believed that as the result of instability cold dense clumps will form in the hot flow if the growth of perturbations is fast enough. This mechanism has been used to explain the condensations of galaxies from the intergalactic medium, formation of solar prominence, and condensations in planetary nebulae (e.g., Field 1965). In the case of accretion flows, we should compare the accretion timescale t_{acc} with the growth timescale of perturbations t_{grow} . For LHAFs, from eq. (8), the timescale of the growth of local perturbations is

$$t_{\text{grow}} = (n\Omega)^{-1} = 3G^{-1} = 3 \left[\frac{Q_{\text{rad}}}{W} \left(\frac{\partial \ln Q_{\text{rad}}}{\partial \ln \Sigma} \right)_W \right]^{-1} = \frac{3W}{2Q_{\text{rad}}} \quad (12)$$

This equals the thermal timescale of accretion flows, t_{th} ,

$$t_{\text{grow}} = \frac{3(\gamma - 1)}{2} t_{\text{th}} = t_{\text{th}}, \quad (13)$$

for adiabatic index $\gamma = 5/3$.

We compare the two timescales, t_{acc} and t_{grow} , by numerical calculations. This requires us to solve the global solution of two-temperature LHAFs. This was done in Y01. The results indicate that depending on the value of $\dot{M}$, there are two types of LHAF. When $\dot{M}_1 \lesssim \dot{M} \lesssim \dot{M}_2$, with $\dot{M}_2$ can be as high as $5\dot{M}_1$, the accretion gas in LHAF is hot throughout the disk; when $\dot{M} \gtrsim \dot{M}_2$, up to $\dot{M}_{\text{Edd}}$, within a certain radius the density of the accretion gas is so high that even the sum of compression work and viscous dissipation can not balance the strong Coulomb cooling. As a result, the hot accretion flow will collapse onto the equatorial plane and form an optically thick cold annulus. While the quantitative results such as the value of $\dot{M}_2$ depend on parameters such as α , the qualitative results should not.

Correspondingly, there are two kinds of results when we compare t_{acc} and t_{grow} , as shown by the two lines in Figure 2. The corresponding two models are taken from Fig. 4 of Y01. The solid line corresponds to $\dot{M} = 0.1\dot{M}_{\text{Edd}} < \dot{M}_2$, so the accretion flow is hot throughout the disk; the dashed line to $\dot{M} = 0.3\dot{M}_{\text{Edd}} > \dot{M}_2$, so the hot accretion flow collapses at a radius and form a cold annulus. We see that when $\dot{M} < \dot{M}_2$, $t_{\text{grow}}/t_{\text{acc}} > 1$, so the perturbations have no time to grow before they are swallowed by the black hole, therefore, the thermal instability will have no effect on the dynamics of LHAFs. When $\dot{M} > \dot{M}_2$, $t_{\text{grow}}/t_{\text{acc}} < 1$ at the transition radius. As a result, some cold dense clumps will form there if LHAFs are thermally unstable at that region. We speculate that once the condensation of the hot phase makes $t_{\text{grow}} \approx t_{\text{acc}}$, the condensation will stop. This constraints will be useful when quantitatively calculating the physical states of clumps such as their filling factor, density and temperature.

In our calculations of t_{acc} and t_{grow} , we considered only the “standard” radiative processes, namely synchrotron, bremsstrahlung, and their Comptonization. If we include the additional Compton cooling due to the soft photons from the cold phase, t_{grow} will decrease ($\dot{M}_1$ and $\dot{M}_2$ will also decrease). We expect that we will still have $t_{\text{grow}} > t_{\text{acc}}$ when $\dot{M} < \dot{M}_2$.

Note that LHAF will not collapse into an optically thick cold disk as the result of *local* perturbation since the perturbation wavelength is smaller than the disk scale-height. The thermal stability of LHAFs against long wavelength perturbations is hard to determine, and it is also not clear how fast the growth of perturbations is compared to the accretion timescale if the flow is unstable. On the other hand, if the filling factor of the cold clumps is large enough, it is possible that these clumps may assemble due to frequent collisions and form a disk-like large scale structure.

Many authors have proposed such cold/hot two-phase accretion flow model for AGNs (e.g., Guilbert & Rees 1988; Ferland & Rees 1988; Kuncic, Celletti & Rees 1997). But almost all these work focus on the thermal state of cold clumps and how the clumps re-radiate the energy they absorb. Krolik (1998) also suggested a two-phase accretion flow model. Different

from our suggested physical mechanism of clumps formation, he suggested that such a two-phase feature is the result of the instability in the radiation-pressure-dominated innermost region of a SSD (see e.g., Gammie 1998).

I thank Shoji Kato, Andrei Beloborodov, and Ramesh Narayan for valuable discussions, and the anonymous referee for helpful comments. This work was supported in part by NASA grant NAG5-10780 and NSF grant AST 0307433.

REFERENCES

- Abramowicz, M.A., Chen, X., Kato, S., Lasota, J.-P., & Regev, O. 1995, *ApJ*, 438, L37 (A95)
- Abramowicz, M.A., Czerny, B., Lasota, J.P., & Szuszkiewicz, E., 1988, *ApJ*, 332, 646
- Abramowicz, M.A., Chen, X.-M., Granath, M., Lasota, J.-P, 1996, *ApJ*, 471, 762
- Björnsson G., Abramowicz M.A., Chen, X., & Lasota J.-P. 1996, *ApJ*, 467, 99
- Chen, X., Abramowicz, M.A., Lasota, J-P, Narayan, R. & Yi, I. 1995, *ApJ*, 443, L61 (C95)
- Esin, A.A., McClintock, J. E., & Narayan, R. 1997, *ApJ*, 489, 865
- Fabian, A.C., & Nulsen P.E.J. 1977, *MNRAS*, 180, 479
- Ferland, G.J., & Rees, M.J. 1988, *ApJ*, 332, 141
- Field, G.B. 1965, *ApJ*, 142, 531
- Gammie, C.F., 1998, *MNRAS*, 297, 929
- Guilbert, P.W., & Rees, M.J. 1988, *MNRAS*, 233, 475
- Kato, S., Abramowicz, M.A., & Chen, X., 1996, *PASJ*, 48, 67
- Kato, S. Fukue, J., & Mineshige, S. 1998, *Black-hole Accretion Disks* (Kyoto University Press, Kyoto)
- Kim, W.T. & Narayan, R. 2003, *ApJ*, in press (astro-ph/0303097)
- Krolik, J.H. 1998, *ApJ*, 498, L13
- Kuncic, Z., Celotti, A., & Rees, M.J. 1997, *MNRAS*, 284, 717
- Kusunose, M. & Mineshige, S. 1996, *ApJ*, 468, 330
- Mathews, W.G., & Bregman J.N., 1978, *ApJ*, 224, 308
- Narayan, R., Mahadevan, R., & Quataert, E. 1998, in “The Theory of Black Hole Accretion Discs”, eds. M.A. Abramowicz, G. Björnsson, and J.E. Pringle (Cambridge University Press)
- Narayan, R. & Medvedev, M. 2001, *ApJ*, 562, L129

- Narayan, R. & Yi, I. 1995, ApJ, 444, 231
- Nulsen, P.E.J. 1986, MNRAS, 221, 377
- Paczynski, B., & Wiita, P. J. 1980, A&A, 88, 23
- Pringle, J.E., 1976, MNRAS, 177, 65
- Shakura, N. I.& Sunyayev, R. A. 1973, A&A, 24, 337 (SSD)
- Shapiro, S.L., Lightman, A.P., & Eardley, D. M., 1976, ApJ, 203, 697 (SLE)
- Yuan, F. 1999, ApJ, 521, L55
- Yuan, F. 2001, MNRAS, 324, 119 (Y01)
- Zakamska, N.L. & Narayan, R. 2003, ApJ, 582, 162
- Zdziarski A. A., et al. 1998, MNRAS, 301, 435




Fig. 1.— The thermal equilibrium curve of various accretion solutions. The accretion rate is in units of $\dot{M}_{\text{Edd}} \equiv 10L_{\text{Edd}}/c^2$ and the units of Σ is g cm^{-2} . The parameters are $M/M_{\odot} = 10$, $\alpha = 0.1$, and $R = 5R_s$. The thin solid lines are for $\xi = 1$, representing ADAF, SLE, SSD, and slim disk. The thick solid, dotted and dashed lines are for $\xi = -1, -0.1$ and -10 , respectively, representing LHAFs. The squares are the location of unity scattering optical depth. The filled circles and triangles are the results obtained from the global solutions of ADAFs and LHAFs, respectively.




Fig. 2.— The ratio of the accretion timescale to the timescale of growth of perturbations as a function of radii for two LHAF solutions presented in Fig. 4 of Y01. The solid (dashed) line is for $\dot{M} = 0.1(0.3)\dot{M}_{\text{Edd}}$. Other parameters are $\alpha = 0.1, \beta = 0.5$.